study. We are grateful to Martin Luff who checked several of our specimens, and supplied information on distribution, Stuart Ball who produced the mapping routine for our figures and Roger Key and Adrian Fowles who read and commented on earlier drafts of the paper.

## References

- Dillwyn, L.W., 1829. Memoranda relating to the coleopterous insects found in the neighbourhood of Swansea. Murray & Rees, Swansea.
- Fowles, A.P., 1990. The Coleoptera of shingle banks on the River Ystwyth, Dyfed. *Entomologist's Rec.J. Var.* 101: 209-221.
- Holmes, P.R., Boyce, D.B. & Reed, D.K., 1990. Trechus rivularis in Wales. Entomologist's mon. Mag. 126: 109.
- , 1991. The Welsh Peatland Invertebrate Survey Preliminary Report: Methodology and study sites. CSD report no. 1125, Nature Conservancy Council, Peterborough.
- Hyman, P.S., 1986. A national review of British Coleoptera. Part 1A. A review of the status of British Coleoptera. ISR report no. 64. Nature Conservancy Council, Peterborough.
- Key, R.S., 1987a. Blethisa multipunctata (L.) (Col.: Carabidae), abundant at Rhos Goch Common, Radnorshire. Entomologist's mon. Mag. 123: 116.
- , 1987b. Radnorshire field meeting 6th 8th June 1986. Coleopterists' Newsletter 26: 17-22.
- Lindroth, C.H., 1974. Coleoptera: Carabidae. *Handbooks for the identification of British insects*, **4**(2). Royal Entomological Society of London.
- Luff, M.L. & Wardle, J., 1991. Trechus rivularis (Gyll.) (Col.: Carabidae) in North-umberland. Entomologist's mon. Mag. 127: 42.
- Miles, P.M., 1960. (No title.) Nature in Wales 6: 97.
- Morgan, I.K., 1988. A summary of interesting beetle records from Carmarthenshire. *Dyfed Invertebrate Group Newsletter* 9: 14-16.
- Shirt, D.B. (ed.), 1987. British Red Data Books: 2. Insects. Nature Conservancy Council.
- Tomlin, J.R. Le B., 1912. The Coleoptera of Glamorgan. Part 1. Trans. Cardiff Naturalists Soc. 45: 41-58.

Aplocnemus pini (Redt.), not nigricornis (F.) (Col.: Melyridae) in Epping Forest, Essex.

In the list of Epping Forest Coleoptera by the late F.D. Buck (1955, Entomologist's mon. Mag. 91: 172-192), the rare A. nigricornis (p. 184) is given as widely distributed in the area, while there is no mention of the less rare A. pini. I have no doubt, however, that this is an error due to the fact that Joy (1932, Pract. Handb. Brit. Beetles, 1: 431) treats our two species as one under the name nigricornis. Actually they are quite distinct, as is now universally recognised (cf., for instance, Allen, 1975, Entomologist's mon. Mag. 111: 210). I have found pini more than once in Epping Forest and also not far away at Waltham Abbey and Cheshunt but never nigricornis, which I have taken only in East Kent (Orlestone Forest, Ham Street).—A.A. ALLEN, 49 Montcalm Road, Charlton, SE7 8QG.